

PRISTINE C₆₀ FULLERENES INHIBIT THE RATE OF TUMOR GROWTH AND METASTASIS

S.V. Prylutska^{1,*}, A.P. Burlaka², Yu.I. Prylutskyi¹, U. Ritter³, P. Scharff³

Joint Ukrainian-German Center on Nanobiotechnology,

¹Taras Shevchenko National University of Kyiv, Institute of Biology,
Volodymyrska Str., 64, 01601 Kyiv, Ukraine

²R.E. Kavetsky Institute of Experimental Pathology, Oncology and Radiobiology of NAS of Ukraine,
Vasylkivska Str. 45, 03022 Kyiv, Ukraine

³Technical University of Ilmenau, Laboratory of Chemistry, 09684 Ilmenau, Germany

Aim: To estimate the impact of C₆₀ fullerene aqueous solution (C₆₀FAS) on the rate of transplanted malignant tumor growth and metastasis. **Methods:** Lewis lung carcinoma was transplanted into C57Bl/6J male mice. Conventional methods for the evaluation of antitumor and antimetastatic effects have been used. **Results:** The C₆₀FAS at low single therapeutic dose of 5 mg/kg inhibited the growth of transplanted malignant tumor (antitumor effect) and metastasis (antimetastatic effect): the maximum therapeutic effect was found to be of 76.5% for the tumor growth inhibition; the increase of animal life span by 22% was found; the metastasis inhibition index was estimated as 48%. **Conclusion:** It was found that water-soluble pristine C₆₀ fullerenes efficiently inhibit the transplanted malignant tumor growth and metastasis. **Key Words:** Water-soluble pristine C₆₀ fullerenes, Lewis lung carcinoma, tumor growth.

Current status of anticancer therapy indicates the necessity for active search of new agents that will be effective against primary tumor and metastases but demonstrate the minimal level of side effects. As promising antitumor agents could be proposed C₆₀ fullerenes — a unique class of carbon allotropes, which exhibits the biological activity both *in vitro* and *in vivo* [1–3].

The water-soluble pristine (unmodified) C₆₀ fullerenes are nontoxic at low physiological concentrations [4–7], they can penetrate through the membrane of cells [8–10] and have strong antioxidant properties [11]. Murugesan et al. [12] have demonstrated the substantial antiangiogenic activity of C₆₀ fullerenes against either basic fibroblast growth factor- or vascular endothelial growth factor-induced angiogenesis in the chick chorioallantoic membrane model. It is known that an imbalance in the levels of these factors causes many serious diseases including malignant growth [13].

Thus, the purpose of this study was to evaluate the impact of water-soluble pristine C₆₀ fullerenes on the transplanted tumor growth and metastasis.

The samples of C₆₀ fullerene aqueous solution (C₆₀FAS) were prepared as follows [14]. We used a saturated solution of pure C₆₀ fullerenes (purity 99.5%) in toluene and the same amount of distilled water in an open beaker. Two phases are formed. Then we applied an ultrasonic bath as long as the toluene needs to evaporate completely. Meanwhile the water phase became yellow colored, indicating that the aqueous fullerene solution has been formed. Thereafter we filtered the aqueous solution from undissolved C₆₀ fullerenes. As a result we prepared the C₆₀FAS sample with maximum concen-

tration of C₆₀ fullerenes in water 1.0 mg/ml. The C₆₀FAS sample is stable during 18 months at 4 °C.

Theoretical calculations [15–16] showed that C₆₀FAS contains both the single C₆₀ molecules and their clusters and solids (with sizes of ~0.7–4 nm in dependence of C₆₀ fullerene concentration in water) in the hydrated state. Moreover, C₆₀ fullerenes structure the water, absorbed by DNA molecules [17], and thus they can affect the DNA functioning in the biological systems.

State of C₆₀ fullerenes in water was monitored using STM technique (NT-MDT, Russia). Samples were deposited on Au(111) surface by precipitation from aqueous solution droplet.

It is important to note that used C₆₀FAS in our experiments does not show a cytotoxic effect with respect to both normal and transformed cells at concentrations below 1.0 mg/ml [6].

The male mice of C57Bl/6J line (20–21 g of b.w.) were kept in a vivarium on a standard diet. The average temperature in a vivarium was 20±1 °C. All experiments were performed in accordance with the international principles of European Convention for protection of vertebrate animals.

Tumor transplantation (Lewis lung carcinoma) was performed by intramuscular injection to the animal's limb (initial number of tumor cells ~5·10⁵, antitumor effect) or to the pad of animal's limb (initial number of tumor cells ~1·10⁶, antimetastatic effect). It is well known that this tumor is characterized by a high degree of metastasis into the lung.

The C₆₀FAS in the volume of 0.1 ml (initial concentration of C₆₀ fullerenes in water was 1.0 mg/ml) was injected intraperitoneally to the animals with transplanted tumor (group 1). Injection of C₆₀FAS was started in a day after transplantation of tumor, which visually appeared on the 10th day. The schedule of C₆₀FAS administration was based on the data obtained by [18].

Received: July 7, 2011.

*Correspondence: E-mail – prylut@ukr.net

Abbreviations used: C₆₀FAS – C₆₀ fullerene aqueous solution

Finally, group 0 (mice with transplanted tumor without C₆₀FAS injection) was used as a control. Initial number of animals in each group was 5 in Experiment 1 (antitumor effect), 7 in Experiment 2 (antitumor effect) and 7 in Experiment 3 (antimetastatic effect). On the 20th day (Experiment 3) all animals were sacrificed and autopsied to calculate the number of metastases in the lung by conventional method.

It is also important to note that these experiments were performed in different time: Experiment 1 — end of spring — summer 2010; Experiment 2 — end of winter — spring 2011; Experiment 3 — end of spring — summer 2011.

Antitumor effectiveness of the applied technology was estimated by following quantitative indicators:

$k_i^{TGI} = [(V_0 - V_1)/V_0] \cdot 100\%$ — tumor growth inhibition (TGI, %); , where V_0 and V_1 are the average values of tumor volume in animals of group 0 (control) and experimental group 1, respectively; $V = (a + b)^3/16$, where a and b are the length and width (in mm) of the tumor site;

$k_i^{IAL} = [(t_1 - t_0)/t_0] \cdot 100\%$ — increasing of animal life (IAL, %); , where t_0 and t_1 are average life span of animals (in days) in group 0 (control) and experimental group 1, respectively;

$k_i^{MII} = [(A_0 \cdot B_0 - A_1 \cdot B_1)/A_0 \cdot B_0] \cdot 100\%$ — metastasis inhibition index (MII, %); , where A_0 and A_1 are frequency of metastasis in the group 0 (control) and experimental group 1, respectively; B_0 and B_1 are average number of metastases in certain organ of animals in group 0 (control) and experimental group 1, respectively.

Statistical analysis of results was performed using STATISTICA software package. On significance of differences was analysed using the parametric (Student's t-test) method [19]. The differences were considered as valid at $p < 0.05$.

The STM images of submonolayer C₆₀ fullerene film deposited from aqueous solution (C₆₀ fullerene concentration in water was 1.0 mg/ml) on Au(111) surface are shown on Figure. They revealed almost random arrangement of C₆₀ fullerene clusters with sizes up to ~2.8 nm (the first stable sphere-like cluster consisting of 13 hydrated C₆₀ fullerenes [15– 16]) (Figure, a). Despite of the mobility of C₆₀ molecules on Au(111) at room temperature we were able to image single C₆₀ molecules (Figure, b).




Figure. a) Large scale STM image of submonolayer C₆₀ fullerene film deposited from aqueous solution (C₆₀ fullerene concentration in water was 1 mg/ml) on Au(111) surface. Some C₆₀ fullerene clusters are aligned along preferential direction <112>; b) Single C₆₀ fullerenes. Lateral size is increased because of shape of the tip. Inset: cross-section along line AB. Scanning parameters: $I_s=40$ pA, $U_s=0.7$ V.

Results of antitumor effect of C₆₀FAS, obtained in Experiment 1 and Experiment 2, are presented in Table 1 and 2, respectively.

Table 1. Experiment 1 (antitumor effect): start of tumor transplantation – 25.05.2010. Start of C₆₀FAS injection after tumor transplantation – 27.05.2010

Days after tumor transplantation	Group 0 (control) n	Group 1 (injection of C60FAS after tumor transplantation)	
		n	Tumor growth inhibition, k_i^{TGI} , %*
11	5	5	54.5
14	5	5	71.9
17	5	5	74.0
21	3	5	74.4
23	2	5	76.5
25	2	5	73.1
28	2	5	
30	1	5	
36	0	5	
39		5	
43		3	
66		1	
77		0	

Note: n – number of mice; *the differences are statistically significant compared with the control ($p < 0.05$).

Table 2. Experiment 2 (antitumor effect): start of tumor transplantation – 17.02.2011. Start of C₆₀FAS injection after tumor transplantation – 19.02.2011. The increase of animal life span (k_i^{IAL} , %) is given in parentheses

Days after tumor transplantation	Group 0 (control) N	Group 1 (injection of C60FAS after tumor transplantation)	
		n	Tumor growth inhibition, k_i^{TGI} , %*
13	7	7	4.0
16	7	7	25.1
21	7	7	5.4
23	4	7	17.0
27	2	6	5.0
29	2	6	
30	0	6	
33		4	
34		2	
37		0	

$(k_i^{IAL}=21.8\%)*$

Note: n – number of mice; *the differences are statistically significant compared with the control ($p < 0.05$).

The inhibition effect of C₆₀ fullerenes on the tumor growth was observed on the 11th day (Experiment 1) and the 13th day (Experiment 2) of the experiment. The maximum value of inhibition of the tumor growth was found to be of 76.5% on the 23th day (Experiment 1) and of 25.1% on the 16th day (Experiment 2) of the experiment. Last animal in the control group 0 died on the 36th day (Experiment 1) and the 30th day (Experiment 2) of the experiment. It should be noted that all 5 mice, injected by C₆₀ fullerenes (group 1; Experiment 1), lived 39 days. Last animal of the experimental group 1 died on the 77th day (Experiment 1) and the 37th day (Experiment 2) of the experiment. The increase of animal life was found to be of 21.8% in Experiment 2. Moreover, it was determined that the average tumor volume in the control group exceeded this parameter in the group 1 by ~3 fold on the 28th day after tumor transplantation in Experiment 2.

The metastasis inhibition index (k_i^{MII}) was obtained as 48% in the study of antimetastatic effect of C₆₀FAS (Experiment 3).

In conclusion, the C₆₀FAS containing hydrated single C₆₀ molecules and C₆₀ clusters with size up to ~2.8 nm without showing direct cytotoxicity at low sin-

gle therapeutic dose of 5 mg/kg [6] demonstrates the inhibition of growth of transplanted tumor (antitumor effect): the maximum therapeutic effect was determined as 76.5% in Experiment 1 and 25.1% in Experiment 2; the increase of animal life span by 21.8% was found in Experiment 2. The metastasis inhibition index was obtained as 48% (antimetastatic effect, Experiment 3). Finally, the anticancer affect of C₆₀FAS was confirmed by histological data for the animal tumors of control and investigated group in Experiment 3 [20]. The obtained results can be explained as a result of the high antioxidant activity of C₆₀ fullerenes [11], neutralizing excess reactive oxygen species in the cell [20], and possibly blocking the specific cell receptors, for example, endothelial growth factor receptors [12]. The proposed method of C₆₀FAS application for the inhibition of tumor growth is promising for experimental oncology and may be tested in the future studies [21].

ACKNOWLEDGMENTS

This work was partly supported by BMBF grant.

REFERENCES

1. Freitas Jr RA. Nanomedicine, Volume IIA: Biocompatibility. Georgetown, TX: Landes Bioscience, 2003.
2. Medicinal chemistry and pharmacological potential of fullerenes and carbon nanotubes. Series: Carbon Materials: Chemistry and Physics. F. Cataldo, T. Da Ros (Eds.). Netherlands: Springer, 2008.
3. Anilkumar P, Lu F, Cao L, *et al.* Fullerenes for applications in biology and medicine. *Current Med Chem* 2011; **18**: 2045–59.
4. Andrievsky G, Klochkov V, Derevyanchenko L. Is the C₆₀ Fullerene Molecule Toxic?! Fullerenes, Nanotubes, and Carbon Nanostruct 2005; **13**: 363–76.
5. Levi N, Hantgan RR, Lively MO, *et al.* C₆₀ fullerenes: detection of intracellular photoluminescence and lack of cytotoxic effects. *J Nanobiotechnol.* 2006; **4**: 4–14.
6. Prylutska SV, Matyshevska OP, Golub AA, *et al.* Study of C₆₀ fullerenes and C₆₀-containing composites cytotoxicity *in vitro*. *Mater Sci Engineer C* 2007; **27**: 1121–4.
7. Johnston HJ, Hutchison GR, Christensen FM, *et al.* The biological mechanisms and physicochemical characteristics responsible for driving fullerene toxicity. *Toxicol Sci* 2010; **114**: 162–82.
8. Foley S, Crowley C, Smaih M, *et al.* Cellular localisation of a water-soluble fullerene derivative. *Biochem Biophys Res Commun* 2002; **294**: 116–9.
9. S.V. Prylutska, O.P. Matyshevska, I.I. Grynyuk, *et al.* Biological effects of C₆₀ fullerenes *in vitro* and in a model system. *Mol Cryst Liq Cryst* 2007; **468**: 265–74.
10. Schuetze C, Ritter U, Scharff P, *et al.* Interaction of N-fluorescein-5-isothiocyanate pyrrolidine-C₆₀ compound with a model bimolecular lipid membrane. *Mater Sci Engineer C* 2011; **31**: 1148–50.
11. Prylutska SV, Grynyuk II, Matyshevska OP, *et al.* Anti-oxidant properties of C₆₀ fullerenes *in vitro*. Fullerenes, Nanotubes, Carbon Nanostruct 2008; **16**: 698–705.
12. Murugesan S, Mousa SA, O'Connor LJ, *et al.* Carbon inhibits vascular endothelial growth factor- and fibroblast growth factor-promoted angiogenesis. *FEBS Lett* 2007; **581**: 1157–60.
13. Carmeliet P. Angiogenesis in life, disease and medicine. *Nature* 2005; **438**: 932–6.
14. Scharff P, Risch K, Carta-Abelmann L, *et al.* Structure of C₆₀ fullerene in water: spectroscopic data. *Carbon* 2004; **42**: 1203–6.
15. Prilutski YuI, Durov SS, Yashchuk VN, *et al.* Theoretical predictions and experimental studies of self-organization C₆₀ nanoparticles in water solution and on the support. *Eur Phys J D* 1999; **9**: 341–3.
16. Bulavin L, Adamenko I, Prylutsky Yu, *et al.* Structure of fullerene C₆₀ in aqueous solution. *Phys Chem Chem Phys* 2000; **2**: 1627–9.
17. Turov VV, Chekhun VF, Krupskaya TV, *et al.* Effect of small addition of C₆₀ fullerenes on the hydrated properties of nanocomposites based on highly dispersed silica and DNA. *Chem Phys Lett* 2010; **496**: 152–6.
18. Gharbi N, Pressac M, Hadchouel M, *et al.* C₆₀ fullerene is a powerful antioxidant *in vivo* with no acute or subacute toxicity. *Nano Lett* 2005; **5**: 2578–85.
19. Gubler EV. Computational methods of analysis and recognition of pathological processes. Moscow: Medicine, 1978, 293 p. (in Russian).
20. Prylutska SV, Burlaka AP, Klymenko PP, *et al.* Using water-soluble fullerenes C₆₀ in anticancer therapy. *Cancer Nanotechnol* 2011; doi: 10.1008/s 12645-011-0020.
21. Jiao F, Qu Y, Zhou G, *et al.* Modulation of oxidative stress by functionalized fullerene materials in the lung tissues of female C57/BL mice with a metastatic Lewis lung carcinoma. *J Nanosci Nanotechnol.* 2010; **10**: 8632–7.22.
22. Prylutska SV, Burlaka AP, Prylutsky YuI. Application of unmodified C₆₀ fullerenes as anticancer agents in the therapy of malignant tumors. Patent of Ukraine for an invention No a201104882, April 14, 2011.