

Cancer Research



Inhibition of Human Breast Cancer Cell Proliferation in Tissue Culture by the Neuroleptic Agents Pimozide and Thioridazine

Jeannine S. Strobl, Keith L. Kirkwood, Tracy K. Lantz, et al.

Cancer Res 1990;50:5399-5405.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/50/17/5399>

- E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.
- Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.
- Permissions** To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.

Inhibition of Human Breast Cancer Cell Proliferation in Tissue Culture by the Neuroleptic Agents Pimozide and Thioridazine¹

Jeannine S. Strobl,² Keith L. Kirkwood, Tracy K. Lantz, Michael A. Lewine, Virginia A. Peterson, and Jennings F. Worley III

Department of Pharmacology and Toxicology, West Virginia University Health Sciences Center, Morgantown, West Virginia 26506

ABSTRACT

Permanent cell culture lines derived from human breast cancer tissue are important experimental models in the study of human breast cancer cell proliferation. In the present work, pimozide, thioridazine, W-13, and W-12 were shown to inhibit MCF-7 human breast cancer cell growth. The 50% inhibition concentration values determined in two proliferation assays, [³H]thymidine incorporation and cell number, were in close agreement for each compound tested. The order of potency for growth inhibition in the presence of 2% stripped calf serum was pimozide (K_i 2 μ M) > thioridazine (K_i 5 μ M) > W-13 (K_i 15 μ M) > W-12 (K_i 39 μ M). Similar concentrations of these compounds blocked estradiol-induced growth of MCF-7 cells, but estrogen receptor (ER) interactions do not seem to be involved. Pimozide and thioridazine had no effect on the estradiol binding properties of the MCF-7 ER, nor did pimozide interfere with the induction of progesterone receptors by estradiol. Furthermore, pimozide also inhibited incorporation of [³H]thymidine into MCF-7 cells stimulated by polypeptide hormones in serum-free medium. The K_i for pimozide in serum-free medium alone, 0.46 μ M, was similar to that determined in the presence of insulin (0.42 μ M), insulin-like growth factor I (0.54 μ M), and epidermal growth factor (0.43 μ M). The effects of pimozide on breast cancer cell growth were not limited to the MCF-7 cell line. Pimozide also blocked cell growth and [³H]thymidine incorporation into the ER-positive T47D and ZR75-1B human breast cancer cell lines and the ER-negative human breast cancer cell line, MDA-MB-231. Although numerous mechanisms of action of pimozide and thioridazine have been identified, both drugs are calmodulin antagonists at drug concentrations that inhibit breast cancer cell growth *in vitro*. Inhibition of MCF-7 cell growth by the selective calmodulin antagonists W-13 and W-12 is consistent with a role for calmodulin antagonism in the broad growth-inhibitory properties of pimozide. We conclude that pimozide and thioridazine may be useful in the control of estradiol- and polypeptide hormone-induced growth of ER-positive and ER-negative human breast tumors.

INTRODUCTION

Estrogens are involved in the etiology and growth of human breast cancer and are thought to exert their effects via binding to specific, high affinity estrogen receptors. The presence of ER³ in human breast cancer is predictive of hormone-dependent tumors which regress in response to endocrine therapies in about 50% of patients (1). The remaining ER-positive and ER-negative tumors are generally resistant to endocrine therapies,

and these are classified as "hormone independent." Alternative therapies for estrogen-dependent tumors and for tumors that are resistant to existing endocrine treatments may be possible based upon an understanding of the mechanisms controlling breast cancer cell proliferation in response to polypeptide hormones. Insulin, insulin-like growth factor I and II, epidermal growth factor, transforming growth factor- α , and basic fibroblast growth factor have been shown to stimulate proliferation of human breast cancer cells *in vitro* (2). Autocrine and paracrine production of these or related polypeptide growth factors may explain the estrogen-independent growth of certain ER-positive and ER-negative human breast cancers (3).

Recent studies of the mechanism of growth regulation by polypeptide hormones in fibroblasts indicate that several intracellular signal transduction pathways are utilized in the transmission of the mitogenic stimulus to the nucleus (4). Various calmodulin antagonists (5-10) and protein kinase C inhibitors (11-14) have been shown to inhibit cell growth *in vitro*, but pharmacological agents with multiple intracellular actions may be particularly effective inhibitors of cell proliferation. Tamoxifen, an antiestrogen used widely in the treatment of ER-positive human breast cancers (15), exerts multiple pharmacological effects in addition to inhibition of estrogen-receptor interactions. These include binding to a distinct antiestrogen binding site (16), receptors for histamine (17) and dopamine (18), inhibition of calmodulin (19) and protein kinase C activation (20), and blockade of voltage-dependent plasma membrane calcium channels (21). High concentrations of tamoxifen inhibit growth of ER-negative human breast cancer cell lines in tissue culture (22). These ER-independent mechanisms may contribute to breast cancer growth inhibition by tamoxifen.

In the experiments described in this report, pimozide inhibited the proliferation of ER-positive and ER-negative human breast cancer cell lines grown in 2% charcoal-stripped calf serum *in vitro*. In MCF-7 cells, pimozide also blocked estrogen-induced growth independently of effects on the ER and proliferation induced by the polypeptide hormones insulin, IGF-I, and EGF. These results in tissue culture were observed with therapeutic drug concentrations, suggesting that pimozide and thioridazine may be of practical value in the management of human breast cancer *in vivo*.

MATERIALS AND METHODS

Cell Culture. Three ER-positive human breast cancer cell lines were used in these studies, MCF-7 (passage 24-69) (23), ZR75-1B (24), T47D (25), and one ER-negative human breast cancer cell line, MDA-MB-231 (26). Cells were grown in DMEM plus 10% FCS (Hazleton Research Products, Inc., Lenexa, KS) and 2 mM glutamine at 37°C in a humidified 5% CO₂-95% air incubator. Experiments were conducted in phenol red-free DMEM (Hazleton Research Products) and 2% stripped CS (GIBCO Laboratories, Grand Island, NY), treated to remove endogenous steroids. To prepare stripped CS, a 0.25% dextran-2.5% Norit A (dextran-coated charcoal) suspension in 0.01 M Tris, pH 7.9, was centrifuged (1,500 \times g, 15 min) to pellet the charcoal. One hundred ml of CS were incubated for 30 min at 56°C with the pellet

Received 2/23/90; revised 5/21/90.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by Pharmaceutical Manufacturer's Association Research Starter Awards to J. S. S. and J. F. W., The Charleston Area Medical Center Foundation, and NIH Grants 3R25CA19530, DK33631, and CA46350.

² To whom requests for reprints should be addressed, at Department of Pharmacology & Toxicology, West Virginia University, Morgantown, WV 26506.

³ The abbreviations used are: ER, estrogen receptor(s); CS, calf serum; DMEM, Dulbecco's modified Eagle's medium; EGF, epidermal growth factor; estradiol, 1,3,5-(10)-estratriene-3,17/ β -diol; FCS, fetal calf serum; HBSS, Hanks' balanced salt solution, 0.4 g/liter KCl-0.06 g/liter KH₂PO₄-8 g/liter NaHCO₃-0.09 g/liter Na₂HPO₄·7H₂O; IGF-I, insulin-like growth factor I; pimozide, 1-(1-(4,4-bis(4-fluorophenyl)butyl)-4-piperidinyl)-1,3-dihydro-2H-benzimidazol-2-one; TED, 0.01 M Tris(hydroxymethyl)aminomethane-1.5 mM disodium EDTA-5 mM dithiothreitol, pH 7.4; thioridazine, 10-[2-(1-methyl-2-piperidyl)ethyl]-2-(methylthio)phenothiazine; W-12, N-(4-aminobutyl)-2-naphthalenesulfonamide hydrochloride; W-13, N-(4-aminobutyl)-5-chloro-2-naphthalenesulfonamide hydrochloride.

from 20 ml of 0.25% dextran-2.5% Norit A suspension. The CS was centrifuged ($1,500 \times g$, 15 min) to remove the charcoal and then incubated for 2 h at 37°C with 2 units/ml of type IV sulfatase from limpets (Sigma, St. Louis, MO). The initial treatment with dextran-Norit A was repeated, followed by a 30-min centrifugation at $12,000 \times g$. The stripped serum was sterilized by filtration through a $0.22\text{-}\mu\text{m}$ pore filter and stored frozen at -20°C until use.

Drugs and Hormones. Pimozide (Lemmon Co., Sellersville, PA) was prepared in dimethyl sulfoxide and stored at -20°C . Thioridazine (Smith, Kline, and Beckman, Philadelphia, PA), W-13, and W-12 (Seikagaku America, Inc., Rockville, MD) were prepared fresh in water for each experiment and sterilized by filtration.

Estradiol- 17β (Steraloids, Pawling, NY) and [2,4,6,7- ^3H]estradiol (91 Ci/mmol; Amersham Corp., Arlington Heights, IL) were stored in 95% ethanol at -20°C . Promegestone (R5020) and [17 α -methyl- ^3H]promegestone (82 Ci/mmol) were purchased from DuPont (Wilmington, DE) and stored in 95% ethanol in the dark at -20°C . Insulin (bovine pancreatic), insulin-like growth factor I (human, recombinant), and epidermal growth factor (mouse submandibular gland) were purchased from Boehringer-Mannheim (Indianapolis, IN) and stored at 4°C as sterile saline solutions.

Cell Growth Determinations. Cells were replica plated ($0.5\text{--}3 \times 10^5$ cells) in T-25 flasks in DMEM plus 10% FCS and 2 mM glutamine. The plating medium was replaced three times at 6–8-h intervals with fresh 3-ml aliquots of phenol red-free DMEM plus 2% stripped CS, 2 mM glutamine, and 50 $\mu\text{g}/\text{ml}$ gentamicin. Drug, estradiol, or the appropriate solvent controls were added to the flasks after a total of 24 h in steroid-free medium. On the days indicated, cells in the medium and those harvested from the monolayer by trypsinization were pooled, pelleted by centrifugation at $1000 \times g$ for 5 min, and resuspended in 1 ml of HBSS. Single cell suspensions were made by passing the cells through a 20-gauge needle. Cell counts were performed using a hemacytometer in the presence of 0.06% trypan blue to assess viability.

[^3H]Thymidine Incorporation Assay. Cells were replica plated ($1\text{--}2 \times 10^5/\text{well}$) in 6-well dishes on day 1 in phenol red-free DMEM plus 2% stripped CS. The medium was replaced with fresh phenol red-free DMEM plus 2% stripped CS twice on day 3 and once on day 4. Drugs and/or hormones were added to the cells following the final medium replacement on day 4. Eighteen to 24 h later, cells were incubated at 37°C for 1 h with $0.5 \mu\text{Ci}/\text{well}$ of [^3H -methyl]thymidine (85 Ci/mmol; DuPont). Cells were then placed on ice, washed three times with HBSS, and harvested using HBSS plus 0.1% EDTA. Cell pellets were sonicated, and aliquots were assayed for [^3H]thymidine incorporation into trichloroacetic acid precipitates, as detailed previously (27).

Flow Cytometry. Cells were replica plated ($4 \times 10^5/60\text{-mm}^2$ dish) on day 1 in 3 ml of phenol red-free DMEM plus 2% stripped CS. Medium replacements and drug/hormone additions were performed according to the schedule used in the [^3H]thymidine incorporation assays. Cells were harvested and counted after 24 or 72 h of drug and/or hormone exposure. Nuclei were prepared and stained with propidium iodide according to the method of Vindeløv (28), except that nuclei were treated with trypsin (0.03 mg/ml) for 25 min at 37°C before proceeding with the ribonuclease and propidium iodide staining steps. The propidium iodide-stained nuclei from 1×10^6 cells were analyzed using a FACScan Instrument (Becton-Dickinson, Mountainview, CA) equipped with the Consort30 DNA software package. The distribution of cells in G_1/G_0 , S, and $G_2\text{-M}$ phases of the cell cycle was determined by the sum of broadened rectangles method.

Estrogen Receptor Measurements. MCF-7 cells were replica plated ($1\text{--}4 \times 10^5/\text{well}$) in 6-well dishes in DMEM plus 10% FCS. Twenty-four h later, the plating medium was replaced with phenol red-free DMEM plus 2% stripped CS for at least 24 h. Immediately before the addition of [^3H]estradiol, 2 ml/well of serum-free, phenol red-free DMEM were added. Whole cell [^3H]estradiol binding assays were performed as detailed previously using sodium hydroxide to solubilize the cells (29). The number and affinity (K_d) of the ER sites were determined by computer Scatchard plot analysis (30). Competition between unlabeled estradiol, pimozide, thioridazine, and [^3H]estradiol for receptor binding was determined in an identical manner using [^3H]estradiol at a concentration of 5×10^{-10} M and the indicated




Fig. 1. Inhibition of MCF-7 proliferation by pimozide and thioridazine. Proliferation was measured as [^3H]thymidine incorporation during a 1-h pulse in cells incubated for 24 h with pimozide (Δ) or thioridazine (\blacktriangle), or as the number of viable cells after a 5-day incubation with pimozide (\circ) or thioridazine (\bullet). Data are expressed as the mean percentage of solvent controls \pm SE (bars) of 2–5 experiments.

concentrations of unlabeled competitors.

Nuclear estrogen receptors were measured using an exchange assay and a single saturating concentration of [^3H]estradiol (31). Crude nuclei (see below) were resuspended in 1.5 ml T buffer, and aliquots were incubated with 10 nM [^3H]estradiol in the presence and absence of 1 μM unlabeled diethylstilbestrol. Exchange reactions were conducted at 30°C for 30 min followed by a 30-min incubation at 37°C . Reactions were terminated by dilution and three washes with 1.5 ml of ice-cold TED buffer. Nuclear bound [^3H]estradiol was quantitated by scintillation counting ethanol extracts of the washed nuclear pellets and corrected for the amount of protein. In all estrogen receptor determinations, competed and uncompeteds were assayed in triplicate.

Progesterone Receptor Measurements. MCF-7 cells were plated at low density ($2\text{--}5 \times 10^5$ cells/T-75 flask), grown for 10 days in phenol red-free DMEM plus 2% stripped CS, and then exposed to estradiol, pimozide, or the combination of pimozide and estradiol for 72 h. Cells were harvested by trypsinization, washed in HBSS, resuspended in 1 ml of ice-cold TED, and lysed using 50 strokes of a tight-fitting pestle in a Dounce homogenizer. Cytosol was prepared by sequential centrifugation at $800 \times g$ for 20 min to pellet a crude nuclear fraction and at $100,000 \times g$ for 40 min. Progesterone receptors in the resulting cytosol were measured using a competitive binding assay (32). A single saturating concentration of [^3H]R5020 (20 nM) in the absence and presence of a 100-fold excess of unlabeled R5020 was used. Both competed and uncompeteds tubes contained a final concentration of 100 nM hydrocortisone (Sigma) to prevent R5020 binding to glucocorticoid receptors. Samples were incubated in triplicate at $0\text{--}4^\circ\text{C}$ for 18 h. Unbound [^3H]R5020 was removed by a 10-min incubation ($0\text{--}4^\circ\text{C}$) with dextran-coated charcoal and centrifugation ($1,000 \times g$, 5 min). Aliquots of the supernatant were assayed for radioactivity and protein content as detailed previously.

RESULTS

Inhibition of Cell Growth by Pimozide and Thioridazine. The inhibition of MCF-7 cell numbers and [^3H]thymidine incorporation by increasing concentrations of pimozide and thioridazine is shown in Fig. 1. Qualitatively identical inhibition curves were observed when MCF-7 cells were treated with pimozide or thioridazine in the presence of 2.5 nM estradiol. Under the culture condition used, estradiol stimulated a 1.7-fold increase in cell numbers and a 4.7-fold increase in [^3H]thymidine incorporation compared with control cells maintained in 2% CS alone. Inhibition constants for pimozide and thioridazine were estimated from these data using the equation

$$K_i = (1/f_b - 1) (\text{Drug conc.})$$

where f_b = fractional drug response (Table 1). There is close agreement between the K_i determined by [3 H]thymidine incorporation assays following acute drug exposure (18–24 h) and cell number data obtained following a 5-day drug exposure. Under all conditions tested, pimozone was twice as potent as thioridazine in the inhibition of MCF-7 cell proliferation *in vitro*. In preliminary studies (data not shown), the potencies of trifluoperazine and fluphenazine were found to be equivalent to that of thioridazine in the inhibition of MCF-7 cell growth.

Pimozone inhibition of estradiol-induced [3 H]thymidine incorporation into MCF-7 cells was not overcome by increasing the concentration of estradiol to 250 nM. Pimozone (2.5 μ M) inhibited [3 H]thymidine incorporation by 56% in cells exposed to 2.5 nM estradiol and by 61% in cells exposed to 250 nM estradiol. Therefore, competition with estradiol for ER binding sites or an increased rate of estradiol metabolism to less active metabolites is unlikely to account for the growth-inhibitory actions of pimozone. Analogous studies with thioridazine were not performed; however, the interactions of both pimozone and thioridazine with ER were further explored.

Effects of Pimozone and Thioridazine on the MCF-7 Estrogen and Progesterone Receptors. Pimozone and thioridazine failed to compete with [3 H]estradiol for MCF-7 ER binding sites at drug concentrations that antagonized the growth of MCF-7 cells (Fig. 2). Competition between unlabeled estradiol and [3 H]estradiol for ER binding is shown as a control. Changes in the ability of the estrogen receptor to bind estradiol following

long-term exposure to pimozone and thioridazine were also examined. Whole cell ER binding assays were performed after exposure of MCF-7 cells to pimozone (1 μ M) or the appropriate solvent (0.1% dimethyl sulfoxide) for 4 h, 24 h, or 5 days. Scatchard plot analyses of the binding data were used to estimate the number of estrogen receptor sites per cell and their affinity for estradiol. The binding data are summarized in Table 2 and indicate that neither estrogen receptor number nor affinity for estradiol was reduced by chronic exposure of MCF-7 cells to pimozone and thioridazine.

Finally, the biological activity of the estrogen receptor in pimozone-treated MCF-7 cells was demonstrated by its nuclear localization and its ability to induce cytosolic progesterone receptors (Table 3). Estrogen receptors in estrogen-deprived and estradiol-treated cells were localized primarily within the nucleus. Levels of cytosolic estrogen receptors in these same cells were less than 10% of the nuclear levels (data not shown). Pimozone treatment did not reduce levels of nuclear ER or increase levels of cytosolic ER.

PgR were undetectable in MCF-7 cells grown in the complete absence of estrogen for 10 days. Induction of PgR by treatment with 2.5 nM estradiol for 72 h occurred in both the absence and the presence of 2.5 μ M pimozone. These data suggest that pimozone interferes with the process of estrogen-induced growth in MCF-7 cells via a mechanism that is distinct from that involved in the induction of progesterone receptors.

Effects of Pimozone on MCF-7 Cell Cycle. To determine the influence of pimozone on progression of MCF-7 cells through the cell cycle, MCF-7 cells were treated for 24 or 72 h with estradiol, pimozone, or estradiol plus pimozone (Fig. 3). Pimozone treatment resulted in a concentration- and time-dependent decrease in the percentage of cells in S phase. Prolonged exposure to pimozone resulted in a gradual accumulation of cells in G_1/G_0 . This effect was concentration dependent and most marked in the estradiol-treated cells that were traversing the cell cycle most rapidly. MCF-7 cells exposed to high concentrations of pimozone (10 μ M) in the absence of estradiol showed some accumulation in G_2-M , an effect that is sometimes indicative of cytotoxicity (33). Cell viability was monitored throughout these experiments by trypan blue exclusion, which corroborated this observation. At 24 h, cell viability exceeded 90% in all cells exposed to 0 or 2.5 μ M pimozone and 85% in cells exposed to 5 or 10 μ M pimozone. By 72 h, cell viability in untreated cells was 88%, but it decreased to approximately 65 and 48% in cells treated with 5 or 10 μ M pimozone, respectively. We conclude that a delay in the G_1/G_0 to S phase transition plays a role in the inhibition of MCF-7 cell growth by concentrations of pimozone $<5 \mu$ M and that higher concentrations of pimozone are cytotoxic.

Inhibition of Polypeptide Hormone-induced Growth of MCF-7 Cells by Pimozone. The inhibition of polypeptide hormone-induced MCF-7 cell proliferation was measured in [3 H]thymidine incorporation assays in which the final medium change contained no serum. The lowest concentrations of insulin, IGF-I, and EGF that stimulated maximal [3 H]thymidine incorporation under these conditions were determined and used in these studies. Parallel MCF-7 cultures were stimulated with insulin, EGF, IGF-I, and estradiol for 18 h in the presence and absence of pimozone. The K_i values for pimozone calculated for all three polypeptide hormones were indistinguishable (Table 4). Pimozone inhibited estradiol-stimulated [3 H]thymidine incorporation in serum-free medium with a K_i of $0.25 \pm 0.12 \mu$ M. The similarity in the inhibition constants for pimozone in unstimulated cells and in cells treated with insulin, IGF-I, EGF, or




Fig. 2. Competition between the thioridazine, pimozone, and estradiol for ER binding in intact MCF-7 cells. Cells were incubated with serum-free phenol red-free DMEM containing 0.5 nM [3 H]estradiol alone or in the presence of the unlabeled competitors at the concentrations indicated for 40 min at 37°C. Specific [3 H]estradiol binding in the absence of competitor is equal to 1. Data represent the mean \pm SE (bars) of 3 experiments performed in triplicate.

Table 2 Estrogen receptors in MCF-7 cells after acute and chronic exposure to pimozone and thioridazine

Time	ER (sites/cell) ^a			
	Control	Pimozone (1 μM)	Control	Thioridazine (2.5 μM)
4 h	165,424 ± 60,000 ^b (K _d = 0.44 ± 0.13 pM) ^c	153,353 ± 28,029 (K _d = 0.56 ± 0.11 pM)	133,568 ± 27,699 (K _d = 0.44 ± 0.13 pM)	142,877 ± 21,760 (K _d = 0.46 ± 0.13 pM)
24 h	191,398 ± 19,767	171,976 ± 46,699 (K _d = 0.39 ± 0.12 pM)	169,965 ± 54,597	202,872 ± 65,985 (K _d = 0.44 ± 0.09 pM)
5 days	120,297 ± 4,541 ^b	137,496 ± 22,480 ^b (K _d = 0.33 ± 0.13 pM)	195,213 ± 80,540 ^b	297,904 ± 61,792 ^b (K _d = 0.37 ± 0.11 pM)

^a Data represent the mean ± SE of 3 separate experiments or mean ± range of 2 experiments.
^b n = 2.
^c The K_d of control cells is the mean of 15 separate experiments that include 4-h, 24-h, and 5-day treatment groups.

Table 3 Effect of pimozone on nuclear estrogen receptor (ER_α) and PgR levels

	ER _α ^a (fm/mg protein)	PgR ^b (fm/mg protein)
Control	80 ± 62	0
Estradiol (2.5 nM)	106 ± 54	52 ± 4
Pimozone (2.5 μM)	90 ± 38	2.5 ± 2.5
Pimozone (2.5 μM) + estradiol (2.5 nM)	104 ± 40	44 ± 15

^a ER_α levels are the mean ± SD of triplicate determinations in 6–7 experiments.
^b PgR levels are the mean ± range of triplicate determinations in 2 experiments.

Table 4 Effect of pimozone on polypeptide hormone-induced [³H]thymidine incorporation into MCF-7 cells in serum-free medium

Hormone	Maximal fold induction of [³ H]thymidine incorporation ^a	K _i pimozone ^b (μM)
None	1.00	0.46 ± 0.02
Insulin (3 ng/ml)	3.20 ± 0.68 (n = 3)	0.42 ± 0.01
IGF-I (0.5 ng/ml)	1.97 ± 0.14 (n = 4)	0.54 ± 0.20
EGF (1 ng/ml)	1.40 ± 0.11 (n = 2)	0.43 ± 0.09

^a Data represent mean ± SE of 3–4 experiments or mean ± range of 2 experiments performed in triplicate.
^b Data represent mean ± SD.




Fig. 3. Cell cycle effects of pimozone in MCF-7 cells. Cells were incubated in phenol red-free DMEM plus 2% stripped CS containing the indicated concentrations of pimozone in the absence (A) or presence (B) of 2.5 nM estradiol for 72 h. The percentage of cells in G₁/G₀ (■), S (●), and G₂-M (▲) phase were determined by flow cytometry. Data represent the mean ± SE (bars) of 2–3 experiments.

estradiol is consistent with a common mechanism of growth inhibition, but further experimentation is needed to address this hypothesis. The K_i values reported in Table 4 are lower than shown in Table 1 and reflect the use of serum-free medium. Inclusion of 2% stripped CS to serum-free medium containing




Fig. 4. Comparison of pimozone effects in ER-positive and ER-negative human breast cancer cell lines. Equal numbers of MCF-7 (Δ), T47D (○), ZR75-1B (●), and MDA-MB-231 (▲) cells were incubated for 18 h with pimozone or solvent alone and then pulse labeled with [³H]thymidine. [³H]Thymidine incorporation is expressed as the mean percentage of control cells ± SE (bars) of 2 (ZR75-1B) or 4 experiments performed in triplicate.

insulin raised the estimated K_i for pimozone approximately 5-fold (data not shown).

Pimozone Inhibition of Proliferation in Other Human Breast Cancer Cell Lines. [³H]Thymidine incorporation assays were performed, directly comparing the effects of pimozone in three ER-positive human breast cancer cell lines (MCF-7, ZR75-1B, and T47D) and one ER-negative cell line (MDA-MB-231) during growth in phenol red-free DMEM plus 2% stripped CS (Fig. 4; Table 5). The response of all three ER-positive breast cancer cell lines to pimozone was very similar. Approximately 5 times more pimozone was required to inhibit thymidine incorporation into the ER-negative cell line, MDA-MB-231, than the most sensitive ER-positive line, T47D. To determine the relative sensitivity of all four human breast cancer cell lines to growth inhibition by pimozone, cell numbers were monitored for 6 days during growth in the presence of 5 μM pimozone. These assays showed pimozone to be an equally effective inhib-

Table 5 Comparison of pimozone inhibition of [³H]thymidine incorporation into ER-positive and ER-negative human breast cancer cell lines

Cell line	K _i pimozone ^a (μM)
MCF-7 ^b	1.72 ± 0.32
ZR75-1B ^b	1.48 ± 0.17
T47D ^b	0.95 ± 0.17
MDA-MB-231 ^c	5.28 ± 1.2 ^d

^a Data represent mean ± SE of determinations in 3–5 experiments.

^b ER-positive cell lines.

^c ER-negative cell line.

^d P < 0.05, Newman-Keuls procedure.




Fig. 5. Inhibition of MCF-7 cell proliferation by W-13. Proliferation was measured as [³H]thymidine incorporation during a 1-h pulse in cells incubated with W-13 for 20 h in the presence (▲) or absence (△) of 2.5 nM estradiol, or as the number of viable cells after a 5-day incubation with W-13 in the presence (●) or absence (○) of 2.5 nM estradiol. Data are expressed as the mean percentage of solvent controls ± SE (bars) of 2 [³H]thymidine experiments and 3–6 cell number experiments.

Table 6 Effects of W-13 and W-12 on MCF-7 cell proliferation assays

	K _i (μM)		
	W-13		W-12,
	[³ H]Thymidine incorporation assay ^a	Cell no. ^b	[³ H]thymidine incorporation assay ^a
Control	15.06 ± 3.6	11.23 ± 2.85	38.93
Estradiol (2.5 nM)	6.31 ± 2.1 ^c	13.5 ± 2.96	ND ^d

^a Data are the mean ± SE of 2 experiments performed in triplicate.

^b Data are the mean ± SE of 3–8 experiments performed in triplicate.

^c P < 0.1, Student's *t* test.

^d Not determined.

itor of proliferation in both the ER-negative and ER-positive cell lines (data not shown).

Inhibition of MCF-7 Cell Proliferation by W-13 and W-12. Inhibition of MCF-7 cell numbers and thymidine incorporation by two relatively selective calmodulin antagonists, W-13 and W-12, was next examined. W-13 and W-12 differ 3.8–5.7-fold in their ability to bind calmodulin and inhibit calmodulin-dependent enzyme activation (5). Comparisons of the relative biological activity of W-13 and W-12 provide a relatively specific method for identification of calmodulin-dependent processes. The results in Fig. 5 show that there is a close correlation between the inhibition of MCF-7 cell numbers after a 5-day exposure to W-13 and inhibition of [³H]thymidine incorporation after a 20-h exposure time. Inhibition curves were essentially identical in control cells and in cells stimulated with 2.5 nM estradiol. Evidence that W-13 is acting through calmodulin is provided in Table 6, which compares the relative activity of W-13 and the dechloro derivative, W-12. The inhibition constants for W-12 (39 μM) and W-13 (6–15 μM) estimated from

these data reflect the expected difference in the ability of these two compounds to inhibit calmodulin.

DISCUSSION

Calmodulin antagonism is likely to play a role in the mechanism of the antiproliferative action of pimozone and thioridazine. Our experiments with pimozone and thioridazine are consistent with earlier studies showing inhibition of the growth of MCF-7 (10, 35, 36) and other tumor cells in tissue culture (5–9) by compounds with calmodulin antagonist activity. Concentrations of pimozone and thioridazine that block breast cancer cell growth are similar to drug concentrations that have been shown to inhibit calmodulin-dependent enzyme activation (34). Furthermore, our results with the naphthalenesulfonamides W-13 and W-12 provide specific evidence for the role of calmodulin in the control of MCF-7 breast cancer cell growth *in vitro*.

Pimozone is the most potent calmodulin antagonist among the neuroleptic drugs currently approved for clinical use (34); thus, we have focused our efforts on this compound. Pimozone was shown to inhibit proliferation of one ER-negative and three ER-positive human breast cancer cell lines. In MCF-7 cells, pimozone was shown to block estradiol-induced growth independently of effects on the ER and to inhibit proliferation in response to three polypeptide hormones, insulin, IGF-I, and EGF.

A variety of compounds possessing anticalmodulin activity induce similar alterations in cell cycle kinetics. Inhibition of MCF-7 cell growth by R24571, tamoxifen, and pimozone is accompanied by a decrease in the number of cells in S phase and a concomitant increase in the number of cells in G₁/G₀ (36, 37). Sutherland and colleagues (36) have presented evidence that the actions of tamoxifen and the calmodulin antagonist R24571 are specifically directed to an event early in the G₁ phase of the MCF-7 cell cycle. These pharmacological data are consistent with a mechanism of action involving a delay in the progression of cells through G₁, and, consistent with this hypothesis, Rasmussen and Means (38) have shown that biochemically elevated intracellular calmodulin concentrations enhance the progression of mouse C127 cells through the G₁ phase of the cell cycle. The nature of the calmodulin-sensitive event(s) in G₁ is of considerable interest, as this could serve as an important target for chemotherapeutic intervention of tumor cell growth.

In light of the evidence that pimozone and thioridazine have calmodulin antagonist activity, it is interesting that no adverse effects on the ability of the MCF-7 ER to bind [³H]estradiol or to function in the induction of PgR were observed in our studies. In calf uterus, there is evidence for calmodulin-dependent phosphorylation of the ER and regulation of estrogen receptor binding (39, 40). In MCF-7 cells, the calmodulin antagonists W-7, R24571, and trifluoperazine have been reported to inhibit the specific uptake of [³H]estradiol and decrease the number of measurable [³H]estradiol-receptor binding complexes in whole cell binding assays (41). These effects disappeared 3 h after exposure to the calmodulin antagonists. Our whole cell binding experiments showed no effect of pimozone or thioridazine on [³H]estradiol binding after 40 min, 4 h, or 5 days of drug exposure at 37°C, and they seem to rule out sustained effects of pimozone and thioridazine on the binding of estradiol to the MCF-7 ER. Moreover, our data are in agreement with Guilino *et al.* (42), who showed that trifluoperazine, W-7, and R24571

failed to compete with estradiol for MCF-7 cytosol ER binding sites.

In addition to calmodulin antagonism, pimozide and thioridazine have multiple cellular actions that may contribute to their antiproliferative effects. Pimozide and thioridazine block calcium entry through plasma membrane calcium channels in tissues such as brain, skeletal muscle, smooth muscle, and pituitary cells (43–46). Although the role of calcium in MCF-7 cell proliferation is not known, a preliminary characterization of calcium channels in MCF-7 cells has been reported (47). Pimozide has also been shown to inhibit protein kinase C activity (48). Although down-regulation of protein kinase C in MCF-7 cells by 12-*O*-tetradecanoylphorbol-13-acetate results in growth stimulation (49), not inhibition, other protein kinases may be important in MCF-7 cell growth regulation. For example, elevations in *c-myc* mRNA levels are associated with estradiol-induced growth in MCF-7 cells (50), and phosphorylation of the *myc*-encoded protein by casein kinase II has been implicated in the transduction of the mitogenic stimuli of insulin, IGF-I, and EGF to the nucleus (51–53).

Pimozide is a neuroleptic that is approved for use in the control of a neurological disorder called Tourette's disease (54). Thioridazine is a neuroleptic drug that is used to control schizophrenia and other psychoses (54). The demonstration that pimozide and thioridazine inhibit the growth of human breast cancer cell lines in tissue culture is evidence that alternative strategies for control of breast cancer proliferation *in vivo* may be found among existing drugs that have not traditionally been considered anticancer agents and provides a rational basis for the experimental use of pimozide *in vivo* to block estrogen- and polypeptide hormone-stimulated breast cancer growth. Significantly, concentrations of pimozide and thioridazine that inhibit breast cancer cell growth *in vitro* are achieved in patients receiving these drugs (43, 55) without the cytotoxic effects associated with standard chemotherapeutic regimens. Control of breast tumor growth *in vivo* by pimozide and thioridazine administration would therefore offer obvious advantages. Some adverse side effects attributable to the antidopaminergic activity of these drugs to occur *in vivo* (54). However, the incidence of Parkinson-like symptoms and tardive dyskinesia with long-term administration of thioridazine is less than observed with other phenothiazines. Elevations in serum prolactin are common in patients receiving all neuroleptic drugs. Although the data implicating prolactin in human breast tumor growth are equivocal (56, 57), prolactin receptors have been identified in 40% of human breast tumors (58). The potential for stimulation of breast tumor growth *in vivo* by pimozide and thioridazine is of concern and could affect the clinical usefulness of pimozide and thioridazine in breast cancer treatment. The concurrent use of a prolactin-lowering drug such as bromocriptine (59) in patients with prolactin receptor-positive tumors may suffice to control prolactin-stimulated tumor growth. We conclude that pimozide and thioridazine are possible candidates for further development for use in human breast cancer treatment.

ACKNOWLEDGMENTS

We thank Diane Bronzert, Karen Huff, and Dr. Michael Moore for cell lines, Cathy Nowak for operation of the FACScan, Dr. Marc Lippman for comments on the manuscript, Cara Williams for assistance with preliminary experiments, and Virginia Lewis and Roxanne Walker for secretarial assistance.

REFERENCES

- Seibert, K., and Lippman, M. Hormone receptors in breast cancer. *Clin. Oncol., J.* 735–794, 1982.
- Karey, K. P., and Sirbasku, D. A. Differential responsiveness of human breast cancer cell lines MCF-7 and T47D to growth factors and 17 β -estradiol. *Cancer Res.,* 48: 4083–4092, 1988.
- Lippman, M. E., Dickson, R. B., Gelmann, E. P., Rosen, N., Knabbe, C., Bates, S., Bronzert, D., Huff, K., and Kasid, A. Growth regulation of human breast carcinoma occurs through regulated growth factor secretion. *J. Cell. Biochem.,* 35: 1–16, 1987.
- Rozengurt, E., Erusalimsky, J., Mehmet, H., Morris, C., Nanberg, E., and Sinnott-Smith, J. Signal transduction in mitogenesis: further evidence for multiple pathways. *Cold Spring Harbor Symp. Quant. Biol.,* 53: 945–954, 1988.
- Hidaka, H., Susaki, Y., Tanaka, T., Endo, T., Ohno, S., Fujii, Y., and Nagata, T. *N*-(6-Aminoethyl)-5-chloronaphthalenesulfonamide, a calmodulin antagonist, inhibits cell proliferation. *Proc. Natl. Acad. Sci. USA,* 78: 4354–4357, 1981.
- Wei, J. W., Hickie, R. A., and Klaassen, D. J. Inhibition of human breast cancer colony formation by anticalmodulin agents: trifluoperazine, W-7, and W-13. *Cancer Chemother. Pharmacol.,* 11: 86–90, 1983.
- Kikuchi, Y., Iwano, I., and Kato, K. Effects of calmodulin antagonists on human ovarian cancer cell proliferation *in vitro*. *Biochem. Biophys. Res. Commun.,* 123: 385–392, 1984.
- Hait, W. N., and Lee, G. L. Characteristics of the cytotoxic effects of the phenothiazine class of calmodulin antagonists. *Biochem. Pharmacol.,* 34: 3973–3978, 1985.
- Matsui, T., Nakao, Y., Kobayashi, N., Koizumi, T., Nakagawa, T., Kishihara, M., and Fujita, T. Effects of calmodulin antagonists and cytochalasins on proliferation and differentiation of human promyelocytic leukemia cell line HL-60. *Cancer Res.,* 45: 311–316, 1985.
- Ford, J. M., Prozialeck, W. C., and Hait, W. N. Structural features determining activity of phenothiazines and related drugs for inhibition of cell growth and reversal of multidrug resistance. *Mol. Pharmacol.,* 35: 105–115, 1989.
- Zinn, K., Keller, A., Whittemore, L. A., and Maniatis, T. 2-Aminopurine selectively inhibits the induction of β -interferon, *c-fos*, and *c-myc* gene expression. *Science (Wash. DC),* 240: 210–213, 1988.
- Meyer, T., Regenass, V., Fabbro, D., Alteri, E., Rosel, J., Muller, M., Caravatti, G., and Matter, A. A derivative of staurosporine (CGP 41251) shows selectivity for protein kinase C inhibition and *in vitro* anti-proliferative as well as *in vivo* anti-tumor activity. *Int. J. Cancer,* 43: 851–856, 1989.
- Matsumoto, H., and Sasaki, Y. Staurosporine, a protein kinase C inhibitor, interferes with proliferation of arterial smooth muscle cells. *Biochem. Biophys. Res. Commun.,* 158: 105–109, 1989.
- Katayama, N., Nishikawa, M., Minami, N., and Shirakawa, S. Putative involvement of protein kinase C in proliferation of human myeloid progenitor cells. *Blood,* 73: 123–130, 1989.
- Jordan, V. C. (ed.). *Estrogen/Antiestrogen Action and Breast Cancer Therapy.* Madison, WI: University of Wisconsin Press, 1986.
- Sutherland, R. L., Murphy, L. C., Foo, M. S., Green, M. D., Whybourne, A., and Krozowski, Z. S. High-affinity anti-estrogen binding site distinct from the oestrogen receptor. *Nature (Lond.),* 288: 273–275, 1980.
- Kroeger, E. A., and Brandes, L. J. Evidence that tamoxifen is a histamine antagonist. *Biochem. Biophys. Res. Commun.,* 131: 750–755, 1985.
- Hiemke, C., and Ghraf, R. Interactions of non-steroidal antiestrogens with dopamine-receptor binding. *J. Steroid Biochem.,* 21: 663–667, 1984.
- Lam, H.-Y. P. Tamoxifen is a calmodulin antagonist in the activation of cAMP phosphodiesterase. *Biochem. Biophys. Res. Commun.,* 118: 27–32, 1984.
- O'Brien, C. A., Liskamp, R. M., Solomon, D. H., and Weinstein, I. B. Inhibition of protein kinase C by tamoxifen. *Cancer Res.,* 45: 2462–2465, 1985.
- Greenberg, D. A., Carpenter, C. L., and Messing, R. O. Calcium channel antagonist properties of the antineoplastic antiestrogen tamoxifen in the PC12 neurosecretory cell line. *Cancer Res.,* 47: 70–74, 1987.
- Taylor, J. S., Blanchard, B., and Zava, D. T. Estrogen receptor-mediated and cytotoxic effects of antiestrogens tamoxifen and 4-hydroxytamoxifen. *Cancer Res.,* 44: 1409–1414, 1984.
- Soule, H. D., Vazquez, J., Long, A., Albert, S., and Brennan, M. A human cell line from a pleural effusion derived from a breast carcinoma. *J. Natl. Cancer Inst.,* 51: 1409–1413, 1973.
- Engel, L. W., Young, N. A., Tralka, T. S., Lippman, M. E., O'Brien, S. J., and Joyce, M. J. Establishment and characterization of three new continuous cell lines derived from human breast carcinomas. *Cancer Res.,* 38: 3352–3364, 1978.
- Keydar, I., Chen, L., Karby, S., Weiss, F. R., Delarea, J., Radu, M., Chaitcik, S., and Brenner, H. J. Establishment and characterization of a cell line of human breast carcinoma origin. *Eur. J. Cancer,* 15: 659–670, 1979.
- Cailleau, R., Young, R., Olive, M., and Reeves, W. J., Jr. Breast tumor cell lines from pleural effusions. *J. Natl. Cancer Inst.,* 53: 661–666, 1974.
- Strobl, J. S., and Lippman, M. E. Prolonged retention of estradiol by human breast cancer cells in tissue culture. *Cancer Res.,* 39: 3319–3327, 1979.
- Vindelov, L. L. Flow microfluorometric analysis of nuclear DNA in cells from solid tumors and cell suspensions. *Virchows Arch. B ZellPathol.,* 24: 227–242, 1977.
- Strobl, J. S., Kasid, A., Huff, K. K., and Lippman, M. E. Kinetic alterations

- in estrogen receptors associated with estrogen receptor processing in human breast cancer cells. *Endocrinology*, *115*: 1116-1124, 1984.
30. Scatchard, G. The attraction of proteins for small molecules. *Ann. NY Acad. Sci.*, *51*: 660-672, 1949.
 31. Clark, J. H., Peck, E. J., Markaverich, B. M., and Densmore, C. L. Basic Principles and Measurement of Steroid Hormone Receptors. *In*: M. R. Hughes, W. T. Schrader, and B. W. O'Malley (eds.), *Hormone Action and Molecular Endocrinology*, pp. 1-47. Houston: Biological Associates, Inc., 1990.
 32. Nardulli, A. M., Greene, G. L., O'Malley, B. W., and Katzenellenbogen, B. S. Regulation of progesterone receptor messenger RNA and protein levels in MCF-7 cells by estradiol: analysis of estrogen's effect on progesterone receptor synthesis and degradation. *Endocrinology*, *122*: 935-944, 1988.
 33. Bontenbal, M., Sieuwerts, A. M., Klijn, J. G. M., Peters, H. A., Krijnen, H. L. J. M., Sonneveld, P., and Foekens, J. A. Effect of hormonal manipulation and doxorubicin administration on cell cycle kinetics of human breast cancer cells. *Br. J. Cancer*, *60*: 688-692, 1989.
 34. Levin, R. M., and Weiss, B. Mechanism by which psychotropic drugs inhibit adenosine cyclic 3',5'-monophosphate phosphodiesterase of brain. *Mol. Pharmacol.*, *12*: 581-589, 1976.
 35. Sutherland, R. L., Watts, C. K. W., Hall, R. E., and Ruenitz, P. C. Mechanisms of growth inhibition by nonsteroidal antiestrogens in human breast cancer cells. *J. Steroid Biochem.*, *27*: 891-897, 1987.
 36. Musgrove, E. A., Wakeling, A. E., and Sutherland, R. L. Points of action of estrogen antagonists and a calmodulin antagonist within the MCF-7 human breast cancer cell cycle. *Cancer Res.*, *49*: 2398-2404, 1989.
 37. Taylor, I. W., Hodson, P. J., Green, M. D., and Sutherland, R. L. Effects of tamoxifen on cell cycle progression of synchronous MCF-7 human mammary carcinoma cells. *Cancer Res.*, *43*: 4007-4010, 1983.
 38. Rasmussen, C. D., and Means, A. R. Calmodulin is involved in regulation of cell proliferation. *EMBO J.*, *6*: 3961-3968, 1987.
 39. Migliaccio, A., Rotondi, A., and Auricchio, F. Calmodulin-stimulated phosphorylation and 17 β -estradiol receptor on tyrosine. *Proc. Natl. Acad. Sci. USA*, *81*: 5921-5925, 1984.
 40. Auricchio, F., Migliaccio, A., DiDomenico, M., and Nola, E. Oestradiol stimulates tyrosine phosphorylation and hormone binding activity of its own receptor in a cell-free system. *EMBO J.*, *6*: 2923-2929, 1987.
 41. Lam, H. Y., and Chan, A. C. Calmodulin antagonists inhibit specific uptake of estradiol by human breast cancer cells. *Proc. Am. Assoc. Cancer Res.*, *29*: A987, 1988.
 42. Guilino, A., Barrera, G., Vacca, A., Farina, A., Ferretti, C., Screpanti, I., Dianzani, M. U., and Frati, L. Calmodulin antagonism and growth-inhibiting activity of triphenylethylene antiestrogens in MCF-7 human breast cancer cells. *Cancer Res.*, *46*: 6274-6278, 1986.
 43. Gould, R. J., Murphy, K. M. M., Reynolds, I. J., and Snyder, S. H. Antischizophrenic drugs of the diphenylbutylpiperidine type act as calcium channel antagonists. *Proc. Natl. Acad. Sci. USA*, *80*: 5122-5125, 1983.
 44. Flaim, S. F., Brannan, M. D., Swigart, S. C., Gleason, M. M., and Muschek, L. D. Neuroleptic drugs attenuate calcium influx and tension development in rabbit thoracic aorta: effects of pimozone, penfluridol, chlorpromazine, and haloperidol. *Proc. Natl. Acad. Sci. USA*, *82*: 1237-1241, 1985.
 45. Galizzi, J.-P., Fosset, M., Romey, G., Laduron, P., and Lazdunski, M. Neuroleptics of the diphenylbutylpiperidine series are potent calcium channel inhibitors. *Proc. Natl. Acad. Sci. USA*, *83*: 7513-7517, 1986.
 46. Enyeart, J. J., Shen, S.-S., and Hinkle, P. M. Pituitary Ca²⁺ channels: blockade by conventional and novel Ca²⁺ antagonists. *Am. J. Physiol.*, *253*: C162-C170, 1987.
 47. Worley, J. F., III, and Strobl, J. S. Voltage-dependent calcium channels in MCF-7 human breast cancer cells and inhibition of cell growth by calcium channel antagonists. *Cancer Chemother. Pharmacol.*, *24*(Suppl. 2): S84, 1989.
 48. Aftab, D. T., and Hait, W. N. A semiautomated 96-well plate assay for protein kinase C. *Anal. Biochem.*, *187*: 84-88, 1990.
 49. Guilband, N., Pichon, M. F., Faye, J. C., Bayard, F., and Valette, A. Modulation of estrogen receptors by phorbol diesters in human breast MCF-7 cell line. *Mol. Cell. Endocrinol.*, *56*: 157-163, 1988.
 50. Dubik, D., and Shiu, R. P. C. Transcriptional regulation of *c-myc* oncogene expression by estrogen in hormone-responsive human breast cancer cells. *J. Biol. Chem.*, *263*: 12705-12708, 1988.
 51. Sommercorn, J., Mulligan, J. A., Lozeman, F. J., and Krebs, E. G. Activation of casein kinase II in response to insulin and to epidermal growth factor. *Proc. Natl. Acad. Sci. USA*, *84*: 8834-8838, 1987.
 52. Klarlund, J. K., and Czech, M. P. Insulin-like growth factor I and insulin rapidly increase casein kinase II activity in BALB/c3T3 fibroblasts. *J. Biol. Chem.*, *263*: 15872-15875, 1988.
 53. Luscher, B., Kuenzel, E. A., Krebs, E. G., and Eisenman, R. W. Myc oncoproteins are phosphorylated by casein kinase II. *EMBO J.*, *8*: 1111-1119, 1989.
 54. Baldessarini, R. J. Drugs and the treatment of psychiatric disorders. *In*: A. G. Gilman, L. S. Goodman, T. W. Rall, and F. Murad (eds.), *The Pharmacological Basis of Therapeutics*, pp. 387-445. New York: MacMillan Publishing Co., 1985.
 55. Murphy, K. M. M., Gould, R. J., Largent, B. L., and Snyder, S. H. A unitary mechanism of calcium antagonist drug action. *Proc. Natl. Acad. Sci. USA*, *80*: 860-864, 1983.
 56. Musey, V. C., Collins, D. C., Musey, P. I., Martino-Saltzman, D., and Preddy, J. R. K. Long-term effect of a first pregnancy on the secretion of prolactin. *N. Engl. J. Med.*, *316*: 229-234, 1987.
 57. Kleinberg, D. L. Prolactin and breast cancer. *N. Engl. J. Med.*, *316*: 269-271, 1987.
 58. Ben-David, M., Kadar, T., Wittliff, J., Biran, S., and Schally, A. Characterization of prolactin receptors and their distribution among American and Israeli women with breast cancer: implications for prediction of hormonal dependency and treatment. *Biomed. Pharmacother.*, *42*: 101-109, 1988.
 59. Dauvois, S., Spinola, P. G., and Labrie, F. Additive inhibitory effects of bromocryptine (CB-154) and medroxyprogesterone acetate (MPA) on dimethylbenz[*a*]anthracene (DMBA)-induced mammary tumors in the rat. *Eur. J. Cancer Clin. Oncol.*, *25*: 891-897, 1989.